arXiv:hep-ex/0008070v2 29 Aug 2000

HADRONIC B DECAYS TO CHARM FROM CLEO

SHELDON STONE

Physics Department, Syracuse University, Syracuse, N. Y. 13244-1130, USA
E-mail: stone@phy.syr.edu

Recent results include the measurement of $B \to D^{(*)}\pi^+\pi^-\pi^-\pi^o$ decays, with rates of ~1.8%, observation of a significant $\omega\pi^-$ substructure that appears to result from the decay of the ρ' resonance, whose mass and width we determine as 1418 MeV and 382 MeV, respectively. Using B to charmonium decays we find that the B^o fraction in $\Upsilon(4S)$ decays is nearly half, that the CP asymmetries in $J/\psi K^\mp$ and $\psi' K^\mp$ are small, and we also report on the first observation of $B \to \eta_c K$.

1 Introduction

Understanding hadronic decays of the B is crucial to insuring that decay modes used for measurement of CP violation truly reflect the underlying quark decay mechanisms expected theoretically. Most of the decay rate is hadronic. The semileptonic branching ratio for e^-, μ^- and τ^- totals ~25%.[1] Currently, measured exclusive branching ratios for hadronic B decays total only a small fraction of the hadronic width. The measured modes for the $\overline{B}^o$, including $D^+(n\pi)^-$, $D^{*+}(n\pi)^-$, where $3 \geq n \geq 1$, $D^{+(*)}D_s^{-(*)}$ and J/ψ exclusive, totals only about 10%.[1] Thus our understanding of hadronic B decay modes is not yet well based in data. It is also interesting to note that the average charged multiplicity in hadronic B^o decays is 5.8±0.1.[2] Since this multiplicity contains contributions from the D^+ or D^{*+} normally present in $\overline{B}^o$ decay, we expect a sizeable, approximately several percent, decay rate into final states with four pions.[3]

2 First Observation of the ρ' in B Decays

We have made the first statistically significant observations of six hadronic B decays shown in Table 1 that result from studying the reaction $B \to D^{(*)}\pi^+\pi^-\pi^-\pi^o$.[4]

Table 1. Measured Branching Ratios

Mode	$\mathcal{B}$ (%)
$\overline{B}^o \to D^{*+}\pi^+\pi^-\pi^-\pi^o$	1.72±0.14±0.24
$\overline{B}^o \to D^{*+}\omega\pi^-$	0.29±0.03±0.04
$\overline{B}^o \to D^{+}\omega\pi^-$	0.28±0.05±0.03
$B^- \to D^{*o}\pi^+\pi^-\pi^-\pi^o$	1.80±0.24±0.25
$B^- \to D^{*o}\omega\pi^-$	0.45±0.10±0.07
$B^- \to D^{o}\omega\pi^-$	0.41±0.07±0.04

A significant fraction of the final state is $D^{(*)}\omega\pi^-$, and there is a low-mass resonant substructure in the $\omega\pi^-$ mass. (See Fig. 1) A simple Breit-Wigner fit assuming a single resonance and no background gives a mass of 1418±26±19 MeV with an intrinsic width of 382±41±32 MeV.

We determine the spin and parity of the $\omega\pi^-$ resonance (denoted A temporarily) by considering the decay sequence $B \to A\ D$; $A \to \omega\pi$ and $\omega \to \pi^+\pi^-\pi^o$.

The angular distributions are shown in Fig. 2. Here θ_A is the angle between the ω direction in the A rest frame and the A direction in the B rest frame; θ_ω is the orientation of the ω decay plane in the ω rest frame, and χ is the angle between the A and ω decay planes.

The data are fit to the expectations for the various J^P assignments. The ω polarization is very clearly transverse ($\sin^2\theta_\omega$) and that infers a 1^- or 2^+ assignment. The $\cos\theta_A$ distribution prefers 1^-, as does the fit to all



Figure 1. (left) The background subtracted $\omega\pi^-$ mass spectrum from $\overline{B}^o \to D^{*+}\omega\pi^-$ decays fit to a simple Breit-Wigner shape. (right) Same for $B \to D\omega\pi^-$ decays (D^o and D^+ are summed).



Figure 2. The angular distribution of θ_A (top-left), θ_ω (top-right) and χ (bottom). The curves show the best fits to the data for for different J^P assignments. The 0^- and 1^+ are almost indistinguishable in $\cos\theta_A$, while the 1^- and 2^+ are indistinguishable in $\cos\theta_\omega$ and χ. The vertical axis gives efficiency corrected events, 104 events are used.

three projections. Thus this state is likely to be the elusive ρ' resonance.[5] These are by far the most accurate and least model dependent measurements of the ρ' parameters. The ρ' dominates the final state. (Thus the branching ratios for the $D^{(*)}\omega\pi^-$ apply also for $D^{(*)}\rho'^-$.)

Heavy quark symmetry predicts equal partial widths for $D^*\rho'$ and $D\rho'$. We measure the relative rates to be 1.06±0.17±0.04 ; consistent within our relatively large errors.

Factorization predicts that the fraction of longitudinal polarization of the D^{*+} is the same as in the related semileptonic decay $B \to D^*\ell^-\bar{\nu}$ at four-momentum transfer q^2 equal to the mass-squared of the ρ'

$$\frac{\Gamma_L\left(B \to D^{*+}\rho'^-\right)}{\Gamma\left(B \to D^{*+}\rho'^-\right)} = \frac{\Gamma_L\left(B \to D^*\ell^-\bar{\nu}\right)}{\Gamma\left(B \to D^*\ell^-\bar{\nu}\right)}\Big|_{q^2=m^2_{\rho'}} . \qquad (1)$$

Our measurement of the D^{*+} polarization is (63±9)%. The model predictions in semileptonic decays for a q^2 of 2 GeV^2, are between 66.9 and 72.6%.[6] Fig. 3 shows the measured polarizations for the $D^{*+}\rho'^-$, the $D^{*+}\rho^-$,[7] and the $D^{*+}D_s^{*-}$ final states. The latter is based on a new measurement using partial reconstruction of the D^{*+}.[8] Thus this prediction of factorization is satisfied.



Figure 3. Measured D^{*+} polarization versus model predictions.

3 New Results from $B \to$ *Charmonium* Decays

3.1 $\frac{f_{oo}}{f_{+-}} = \frac{\mathcal{B}\left(\Upsilon(4S)\to B^o\overline{B}^o\right)}{\mathcal{B}\left(\Upsilon(4S)\to B^+B^-\right)}$

We measured an important although often overlooked, parameter that effects every B^o or B^+ branching ratio measured on the $\Upsilon(4S)$ resonance. We determined the ratio of the number of $B^o\overline{B}^o$ pairs versus the number of B^-B^+ pairs by measuring the ratio of widths of two-body final states of the type $\psi^{(')}K^{(*)}$. Since the ψ is I=0 and the B and K are I=1/2, the decay is pure ΔI=1/2 and therefore the individual channels, such as ψK, must have the same decay widths for B^o and B^+. We find $\frac{f_{oo}}{f_{+-}} = 1.04 \pm 0.07 \pm 0.04$, which yields nearly equal rates for charged and neutral B's, i.e. $f_{oo} = 0.49 \pm 0.02 \pm 0.01$.[9]

3.2 Limits on CP Violation in $B^\mp \to J/\psi$ *(or* ψ'*)* $K^\mp$

In the Standard Model only one decay amplitude dominates for decays of the type $B^\mp \to$ *charmonium* $K^\mp$. CLEO measured asymmetries consistent with zero for the J/ψ and ψ' cases: (1.8±4.3± 0.4)% and (2.0±9.1±1.0)%, respectively.[10]

3.3 First Measurement of $\mathcal{B}(B \to \eta_c K)$

We observe exclusive final states with the $c\bar{c}$ being the η_c.[11] The rates are $\mathcal{B}(B^- \to \eta_c K^-) = (6.9^{+2.6}_{-2.1} \pm 0.8 \pm 2.0) \times 10^{-3}$ and $\mathcal{B}(B^o \to \eta_c K^o) = (10.9^{+5.5}_{-4.2} \pm 1.2 \pm 3.1) \times 10^{-3}$. These rates are large, similar to those for the $J/\psi K$ final state.

4 Conclusions

Recent CLEO data has vastly increased our knowledge about hadronic B decays. We have seen the elusive ρ' and have the best measurement of its mass and width. Factorization when applied to polarizations appears to work in D^{*+} plus ρ, ρ' and D_s^* decays. The rates of the $\Upsilon(4S)$ into charged and neutral B's has been accurately determined and the first observation of $B \to \eta_c K$ decays has been made. Much however remains to be learned.

Acknowledgments

We thank N. Isgur, J. Rosner and J. Schechter for useful discussions. We thank the National Science Foundation for support.

References

1. C. Caso *et al.*, *The European Physical Journal* **C3** 1(1998).
2. G. Brandenburg *et al.* (CLEO), *Phys. Rev.* **D61**, 072002(2000).
3. ARGUS previously reported a signal of 28±10 events in $D^{*+}\pi^+\pi^-\pi^-\pi^o$, giving a branching ratio of (3.4±1.8)%. H. Albrecht *et al.*, *Z. Phys.* **C 48**, 543 (1990).
4. M. Artuso *et al.* [hep-ex/0006018].
5. A. B. Clegg and A. Donnachie, *Z. Phys.* **C 62**, 455 (1994).
6. N. Isgur and M. B. Wise, *Phys. Lett.* **B237**, 527 (1990); M. Wirbel *et al.*, *Z. Phys.* **C29**, 627 (1985); N. Neubert, *Phys. Lett.* **B264**, 455 (1991).
7. M. Artuso *et al.* (CLEO), *Phys. Rev. Lett.* **82**, 3020 (1999).
8. S. Ahmed *et al.*, [hep-ex/0008015].
9. J.P. Alexander *et al.*, [hep-ex/0006002].
10. G. Bonvicini *et al.*, [hep-ex/0003004].
11. K.W. Edwards *et al.*, [hep-ex/0007012].